

MINIMAL LEPTON FLAVOUR VIOLATION AND LEPTOGENESIS

V. PORRETTI

*Departament de Física Teòrica, Universitat de València, E-46100, Burjassot, Spain and
Dipartimento di Fisica ‘E. Amaldi’, Università di Roma Tre and INFN, Sezione di Roma Tre,
I-00146 Rome, Italy*

We study the role of leptogenesis in the framework of minimal lepton flavour violation.

Keywords: minimal flavour violation; resonant leptogenesis; neutrinos; mu to e gamma.

1. Outline

If neutrino masses are generated by a see-saw mechanism with heavy Majorana neutrinos, the hypothesis of minimal flavour violation (MFV) in the lepton sector cannot be a simple replica of MFV in the quark sector (sec.2): more parameters are involved (sec.3) and high energy processes like leptogenesis (sec.4) can have a role in the predictions for low energy experiments (sec.5). We conclude summarizing the main points in sec.6.

The introduction in sec.2 is mainly based on Refs.^{1,2}, the rest on Ref.³. Aspects of MFV in the lepton sector have been discussed also in Refs.^{4,5,6,7}.

2. A definition of minimal lepton flavour violation

In the Standard Model (SM) flavour changing neutral currents are absent at tree level and mixing between different families is suppressed by small fermion masses and small CKM angles. Any new physics (NP) that fails to reproduce these special features of the SM is likely to give visible contributions to flavour changing rates. Experiments indicate that this is not the case, so that or flavour violating NP intervenes at a very high energy scale or its flavour structure is somehow hidden behind the SM one. The second option is more appealing. It looks more compatible with the hierarchy problem and offers realis-

tic chances to discover NP directly. A natural and general proposal in this direction is the MFV hypothesis, where the Yukawa matrices are assumed to be the only sources of flavour violation also beyond the SM.

More precisely, we assume that there is flavour violating NP at some high energy scale Λ_{FV} which enters the SM lagrangian with non-renormalizable operators suppressed by inverse powers of Λ_{FV} . In the limit of vanishing Yukawa couplings, the Standard Model lagrangian is invariant under the flavour symmetry group $\mathcal{G} = SU(3)_{QL} \times SU(3)_{uR} \times SU(3)_{dR} \times SU(3)_{LL} \times SU(3)_{eR}$. The invariance of the SM Lagrangian under $\mathcal{G}$ can be formally recovered elevating the Yukawa matrices to spurion fields with appropriate transformation properties under $\mathcal{G}$. The hypothesis of MFV states that this is sufficient to make also NP operators invariant under $\mathcal{G}$.

At the moment, the main motivation for extending the hypothesis of MFV to leptons is the analogy with the quark sector, where it is supported by a large amount of experimental data. We would like to understand whether MFV is a general principle that acts also on leptons. The exact analogy would imply that neutrinos are Dirac particles. In this case, the exceptional smallness of neutrino masses would remain unexplained and flavour changing rates like $\mu \rightarrow e\gamma$ and $\mu - e$ conversion in nuclei would be safely below

current and future experimental limits.

For these reasons, it is more interesting to include in the MFV hypothesis lepton number violation (LNV), that explains the smallness of neutrino masses with the high scale of LNV, $\Lambda_{LNV} \gg \Lambda_{FV}$. Once LNV is introduced, the concept of analogy with quarks is ambiguous. A possible choice is to assume that right-handed neutrinos exist (the counterpart of right-handed up-quarks) and that their Majorana mass matrix is proportional to the identity matrix, so that the only flavour non-diagonal objects are the Yukawa couplings, like in the quark sector. However this does not automatically guarantee that flavour changing rates are completely specified in terms of the masses and mixings measured in experiments, as it happens with quarks.

3. Parameter counting

The basic flavour changing unit we obtain from the spurion analysis of NP operators is $\lambda_\nu^\dagger \lambda_\nu$. In the quark sector the analogue quantity $\lambda_u^\dagger \lambda_u$ can be expressed in terms of quark masses and CKM angles, so that the only unknown in the predictions for flavour changing branching ratios is the overall factor $1/\Lambda_{FV}^2$, coming from the power suppression of NP operators of dimension > 4 :

$$BR(\text{quark MFV}) \sim \frac{f(m_u, m_d, V_{CKM})}{\Lambda_{FV}^4} \quad (1)$$

Here instead we can obtain $\lambda_\nu^\dagger \lambda_\nu$ extracting λ_ν from the see-saw relation

$$m_\nu^{eff} \equiv v^2 \lambda_\nu^T M_R^{-1} \lambda_\nu = U_{PMNS}^* m_{\text{diag}} U_{PMNS}^\dagger \quad (2)$$

written in the basis where the charged lepton Yukawa matrix λ_e and the heavy Majorana neutrinos mass matrix M_R are diagonal. We find

$$\lambda_\nu = \frac{M_R^{1/2}}{v} O H(\phi_1, \phi_2, \phi_3) m_{\text{diag}}^{1/2} U_{PMNS}^\dagger \quad (3)$$

where O is a real orthogonal matrix which depends on three real parameters and H a complex hermitian matrix which also depends on three real parameters $\phi_{1,2,3}$; $H \rightarrow I$ in the

CP limit⁸. In our definition of MFV, right-handed neutrinos are degenerate in mass so that M_R in (3) is a number that we identify with Λ_{LNV} and the matrix O can be rotated away (the Lagrangian is $O(3)$ invariant). Using eq.(3), the basic unit that takes part into low energy flavour changing rates in the MFV framework can be written as

$$\lambda_\nu^\dagger \lambda_\nu = \frac{M_R}{v^2} U_{PMNS} m_{\text{diag}}^{1/2} H^2 m_{\text{diag}}^{1/2} U_{PMNS}^\dagger. \quad (4)$$

Let us count the parameters in eq.(4): there are 9 parameters in principle measurable at low energy (MNS angles, Dirac and Majorana phases, neutrino masses) and 4 unknown parameters: the normalization M_R and $\phi_{1,2,3}$ in the matrix H , which disappear in the see-saw relation (2). With non degenerate right-handed neutrinos we would have 5 unknown parameters more (the 2 mass splittings of M_R and the 3 angles in O).

Assuming that all the baryon asymmetry η_B of the universe has been generated through sphaleron effects by a lepton asymmetry, we can use the observed value of $\eta_B = (6.3 \pm 0.3) \times 10^{-10}$ to get some information on the high energy parameters M_R and $\phi_{1,2,3}$. In fact, at first order leptogenesis depends on the combination

$$\lambda_\nu \lambda_\nu^\dagger = \frac{M_R}{v^2} H m_{\text{diag}} H. \quad (5)$$

4. Analysis of leptogenesis

A necessary condition for generating a lepton asymmetry is the non-degeneracy of heavy neutrinos. We in general expect that the tree-level degeneracy of heavy neutrinos is lifted by radiative corrections. In MFV models the most general form of the allowed mass-splittings is

$$\begin{aligned} \frac{\Delta M_R}{M_R} = & c_\nu \left[\lambda_\nu \lambda_\nu^\dagger + (\lambda_\nu \lambda_\nu^\dagger)^T \right] \\ & + c_{\nu\nu}^{(1)} \left[\lambda_\nu \lambda_\nu^\dagger \lambda_\nu \lambda_\nu^\dagger + (\lambda_\nu \lambda_\nu^\dagger \lambda_\nu \lambda_\nu^\dagger)^T \right] \\ & + c_{\nu\nu}^{(2)} \left[\lambda_\nu \lambda_\nu^\dagger (\lambda_\nu \lambda_\nu^\dagger)^T \right] + c_{\nu\nu}^{(3)} \left[(\lambda_\nu \lambda_\nu^\dagger)^T \lambda_\nu \lambda_\nu^\dagger \right] \\ & + c_{\nu l} \left[\lambda_\nu \lambda_\nu^\dagger \lambda_e \lambda_e^\dagger + (\lambda_\nu \lambda_\nu^\dagger \lambda_e \lambda_e^\dagger)^T \right] + \dots \quad (6) \end{aligned}$$

In a model-independent approach we cannot fix the coefficients in (6) but in a perturbative scenario we expect $c_\nu \sim g_{\text{eff}}/4\pi$ and

$c_{\nu\nu}^{(i)}, c_{\nu l} \sim g_{\text{eff}}^2/(4\pi)^2$. From eq.(6) we can derive some general properties of leptogenesis in MFV models: (i) the term proportional to c_ν does not generate an asymmetry by itself, but (ii) sets the order of magnitude of the mass splitting and naturally gives the condition for resonant leptogenesis: the mass splitting of right-handed neutrinos is comparable to the decay width,

$$\Delta M_{R_{ij}} \sim \Gamma_j = M_R \frac{|\lambda_\nu \lambda_\nu^\dagger|_{jj}}{8\pi}. \quad (7)$$

(iii) The right amount of leptogenesis can be generated even with $\lambda_e = 0$, if all the three parameters $\phi_{1,2,3} \neq 0$. (iv) Since $\lambda_\nu \sim \sqrt{M_R}$, for low values of M_R ($\lesssim 10^{12}$ GeV) the asymmetry generated by the $c_{\nu l}$ term dominates but is typically too small to match the observed value of η_B . In this regime we find the flat dependence on M_R typical of resonant leptogenesis. At $M_R \gtrsim 10^{12}$ GeV the quadratic terms $c_{\nu\nu}^{(i)}$ dominate the generation of the asymmetry, which grows linearly with M_R . These specific features of resonant leptogenesis in MFV can be derived with a general analysis of CP-invariants and reproduced analytically. Properties (i),(iv) ex-




Fig. 1. Baryon asymmetry η_B as a function of the right-handed neutrinos mass M_R for $c_{\nu l} = 0$ (cyan circles) and $c_{\nu l} \neq 0$ (blue crosses).

plain the characteristic behaviour of η_B as function of M_R , shown in fig. 1 for reference values of the parameters. Varying all the parameters in a wide range we find the




Fig. 2. Baryon asymmetry η_B as a function of the right-handed neutrinos mass M_R . The experimental value in log-scale is 9.20 ± 0.02 ; the larger error band takes into account the theoretical uncertainties of a model-independent analysis, in particular the conversion factor of the leptonic asymmetry into the baryonic one, which depends on the field content of the model.

result in fig. 2. In deriving this result we used the analytic formulae for leptogenesis without flavour effects of Ref.⁹ and assumed a loop hierarchy between the coefficients of the mass splittings. Under these assumptions, the right size for η_B can be reached for $M_R \gtrsim 10^{12}$ GeV. The regime $M_R \gg 10^{12}$ GeV is particularly interesting since in this case the CP-violating parameters ϕ_i are very small and we recover the predictive scheme of Ref.².

A MFV model is for instance the Minimal Supersymmetric Standard Model with degenerate right-handed neutrinos at the GUT scale. This scenario has been recently analyzed in Ref.⁷. The same behaviour shown in fig. 2 was found, but the use of a different analysis of leptogenesis¹⁰ and the inclusion of flavour effects slightly enhanced the average value of η_B in the low M_R region so that successful leptogenesis could be achieved for some points of the parameter space down to 10^6 GeV.

5. Effect of high energy parameters on $l_i \rightarrow l_j \gamma$

In the MLFV framework, the effective Lagrangian relevant for the radiative decays $l_i \rightarrow l_j \gamma$ is

$$\mathcal{L}_{\text{eff}} = \frac{1}{\Lambda_{\text{LFV}}^2} \left(c_{RL}^{(1)} O_{RL}^{(1)} + c_{RL}^{(2)} O_{RL}^{(2)} \right), \quad (8)$$

where

$$\begin{aligned} O_{RL}^{(1)} &= g' H^\dagger \bar{e}_R \sigma^{\mu\nu} \lambda_e \Delta_{FCNC} L_L B_{\mu\nu}, \\ O_{RL}^{(2)} &= g H^\dagger \bar{e}_R \sigma^{\mu\nu} \tau^a \lambda_e \Delta_{FCNC} L_L W_{\mu\nu}^a, \end{aligned} \quad (9)$$

and g' (g) and $B_{\mu\nu}$ ($W_{\mu\nu}^a$) are the coupling constant and the field strength tensor of the $U(1)_Y$ ($SU(2)_L$) gauge group. This effective Lagrangian leads to

$$\begin{aligned} BR_{l_i \rightarrow l_j \gamma} &\equiv \frac{\Gamma(l_i \rightarrow l_j \gamma)}{\Gamma(l_i \rightarrow l_j \nu_i \bar{\nu}_j)} \\ &= 384 \pi^2 e^2 \frac{v^4}{\Lambda_{\text{LFV}}^4} \left| \left(\lambda_\nu^\dagger \lambda_\nu \right)_{ij} \right|^2 |c_{RL}^{(2)} - c_{RL}^{(1)}|^2 \\ &\sim \left(\frac{\Lambda_{\text{LNV}}}{\Lambda_{\text{LFV}}^2} \right)^2 f(m_\nu^{\text{eff}}, m_l, U_{\text{PMNS}}, \phi_i) \end{aligned} \quad (10)$$

to be compared with eq.(1), which represents a generic branching ratio in MFV in the quark sector. An important difference is that the ϕ_i parameters, non-measurable in low energy experiments, enter the branching ratio and typically produce an enhancement. For large values of M_R their effect is moderate and MFV predicts $BR(\mu \rightarrow e \gamma)/BR(\tau \rightarrow \mu \gamma) \lesssim 1$. A second crucial difference between eqs.(1) and (10) is the absolute normalization of the branching ratios, that now depends on both the scales of lepton number and lepton flavour violation. We cannot interpret the experimental bounds on $l_i \rightarrow l_j \gamma$ as lower limits on the scale Λ_{LFV} , without independent information on $\Lambda_{\text{LNV}} \equiv M_R$. Leptogenesis provided this information, since the observed value of the baryon asymmetry is most naturally reproduced for $M_R \gtrsim 10^{12}$ GeV. For Λ_{LFV} around the TeV, this enhances the branching ratio for $\mu \rightarrow e \gamma$ up to values in the reach of the MEG experiment.

6. Summary

In this talk we studied leptogenesis in a generic MFV scenario with right-handed Majorana neutrinos degenerate in mass. Radiative corrections lift the tree-level degeneracy of right-handed neutrinos and induce mass-splittings proportional to the neutrino and charged lepton Yukawa couplings. We showed that leptogenesis is viable and most efficient at high values of right-handed neutrino masses ($\gtrsim 10^{12}$ GeV), where it is driven by the mass-splittings quartic in the neutrino Yukawa couplings. As a consequence, predictions for $\mu \rightarrow e \gamma$ are enhanced and should be observable in next experiments for natural values of the scale of NP. High energy CP-violating parameters, that disappear in the see-saw relation but take part into leptogenesis, have a significative impact on low-energy processes. The expectation $BR(\mu \rightarrow e \gamma)/BR(\tau \rightarrow \mu \gamma) \ll 1$, valid in the CP limit, is recovered in the regime of very heavy right-handed neutrinos ($M_R \gg 10^{12}$ GeV).

References

1. G. D'Ambrosio, G. F. Giudice, G. Isidori and A. Strumia, Nucl. Phys. B **645** (2002) 155
2. V. Cirigliano, B. Grinstein, G. Isidori and M. B. Wise, Nucl. Phys. B **728** (2005) 121 [arXiv:hep-ph/0507001].
3. V. Cirigliano, G. Isidori and V. Porretti, arXiv:hep-ph/0607068.
4. V. Cirigliano and B. Grinstein, Nucl. Phys. B **752** (2006) 18 [arXiv:hep-ph/0601111].
5. S. Davidson and F. Palorini, arXiv:hep-ph/0607329.
6. B. Grinstein, V. Cirigliano, G. Isidori and M. B. Wise, arXiv:hep-ph/0608123.
7. G. C. Branco, A. J. Buras, S. Jager, S. Uhlig and A. Weiler, arXiv:hep-ph/0609067.
8. J. A. Casas and A. Ibarra, Nucl. Phys. B **618**, 171 (2001) [hep-ph/0103065]; S. Pascoli, S. T. Petcov and C. E. Yaguna, Phys. Lett. B **564**, 241 (2003) [hep-ph/0301095].
9. S. Blanchet and P. Di Bari, arXiv:hep-ph/0603107.
10. A. Pilaftsis and T. E. J. Underwood, Phys. Rev. D **72**, 113001 (2005); T. E. J. Underwood, arXiv:hep-ph/0605232.